South Australian Murray Darling Basin
Lower Murray Darling
Lachlan
Central West
Murrumbidgee
Hawkesbury–Nepean
Murray
ACT
Mallee
Goulburn Broken
North Central
Wimmera
North East
South East
Southern Rivers
East Gippsland
Port Phillip and Western Port
Corangamite
Glenelg Hopkins
West Gippsland
North West
North

Wentworth
Mildura
Renmark
Paringa
Berri
Red Cliffs
Griffith
West Wyalong
Cowra
Young
Hay
Leeton
Cootamundra
Balranald
Narrandera
Goulburn
Ouyen
Wagga Wagga
Pinnaroo
Swan Hill
Tumut
Canberra
Queanbeyan
Deniliquin
Echuca
Corowa
Albury-Wodonga
Cooma
Warracknabeal
Kyabram
Shepparton
Wangaratta
Bordertown
Nhill
Benalla
Horsham
Bendigo
Naracoorte
Maryborough
Penola
Ararat
Ballarat
Sunbury
Melton
Hamilton
Mount Gambier
Melbourne
Bairnsdale
Lakes Entrance
Cranbourne
Geelong
Sale
Moe
Traralgon
Portland
Warrnambool
Colac
Wonthaggi

Number of threatened ecological communitie s affected by feral pigs (*Sus scrofa*) in the pig's current range

This map shows the number of threatened ecological communities impacted on, or potentially impacted on, by feral pigs within their current range. The current range of pigs is identified in 'Assessing invasive animals in Australia 2008' report while information on threatened species was sourced from the Department's Species Profile and Threats database.

No. Threatened Communities

- 0
- 1
- 2
- 3
- Locality
- NRM Regions
- National Reserve System
- Ramsar Sites

N

0 20 40 80 Km

Datasets sources:

National Vertebrate Pest Assessment 2006 Dataset,
© National Land and Water Audit, 2006.

Ramsar wetlands in Australia
© Australian Government, Department of the Environment, Water, Heritage and the Arts, 2009.

Collaborative Australian Protected Areas Database
© Australian Government, Department of the Environment, Water, Heritage and the Arts, 2006.

Natural Resource Management (NRM) Regions (2009)
© Australian Government, Department of the Environment, Water, Heritage and the Arts, 2009.

Species and Communities of National Environmental Significance Database
© Australian Government, Department of the Environment, Water, Heritage and the Arts, 2009.

Australia, Topographic Data - State Borders 1:10 million,
© Geoscience Australia, 1989.

Produced by:
Environmental Resources Information Network
Australian Government
Department of the Environment, Water, Heritage and the Arts
February 2010.

Datum: Geocentric Datum of Australia 1994.

Caveat:
The information in this map has been compiled from a range of sources. While every effort has been made to ensure accuracy and completeness, no guarantee is given, nor responsibility taken by the ommonwealth for errors or omissions, and the Commonwealth does not accept responsibility in respect of any information or advice given in relation to, or as a consequence of, anything containing herein. Some misalignment may occur between layers due to different scales of data capture.

Australian Government
Department of the Environment, Water, Heritage and the Arts

Threatened Ecological Communities affected or potentially affected by feral pigs:

1. Grassy White Box Woodlands
2. Semi-evergreen vine thickets of the Brigalow Belt (North and South) and Nandewar Bioregions
3. Temperate Highland Peat Swamps on Sandstone
4. White Box-Yellow Box-Blakely''s Red Gum Grassy Woodland and Derived Native Grassland