

Information Sheet on Ramsar Wetlands (RIS)

Categories approved by Recommendation 4.7 (1990), as amended by Resolution VIII.13 of the 8th Conference of the Contracting Parties (2002) and Resolutions IX.1 Annex B, IX.6, IX.21 and IX.22 of the 9th Conference of the Contracting Parties (2005).

This Ramsar Information Sheet has been converted to meet the 2009 – 2012 format, but the RIS content has not been updated in this conversion. The new format seeks some additional information which could not yet be included. This information will be added when future updates of this Ramsar Information Sheet are completed. Until then, notes on any changes in the ecological character of the Ramsar site may be obtained from the Ecological Character Description (if completed) and other relevant sources.

1. Name and address of the compiler of this form:

J G Blackman
Environmental Protection Agency
Northern Regional Centre
Marlow Street
PALLARENDA QLD 4810

FOR OFFICE USE ONLY.

DD MM YY

--	--	--

Designation date

--	--	--	--	--	--

Site Reference Number

Ph: (07) 4722 5270
Fax: (07) 4722 5274

2. Date this sheet was completed/updated:
1999

3. Country:
Australia

4. Name of the Ramsar site:

The precise name of the designated site in one of the three official languages (English, French or Spanish) of the Convention. Alternative names, including in local language(s), should be given in parentheses after the precise name.

Bowling Green Bay, Queensland

5. Designation of new Ramsar site or update of existing site:

Bowling Green Bay, Queensland was designated on 22 October 1993
The previous RIS was dated 16th August 1991.

This RIS is for (tick one box only):

- a) Designation of a new Ramsar site ☐; or
b) Updated information on an existing Ramsar site ☒

6. For RIS updates only, changes to the site since its designation or earlier update:

a) Site boundary and area

The Ramsar site boundary and site area are unchanged: ☐

or

If the site boundary has changed:

- i) the boundary has been delineated more accurately ☐; or
- ii) the boundary has been extended ☐; or
- iii) the boundary has been restricted** ☐

and/or

If the site area has changed:

- i) the area has been measured more accurately ☐; or
- ii) the area has been extended ☐; or
- iii) the area has been reduced** ☐

**** Important note:** If the boundary and/or area of the designated site is being restricted/reduced, the Contracting Party should have followed the procedures established by the Conference of the Parties in the Annex to COP9 Resolution IX.6 and provided a report in line with paragraph 28 of that Annex, prior to the submission of an updated RIS.

b) Describe briefly any major changes to the ecological character of the Ramsar site, including in the application of the Criteria, since the previous RIS for the site:

7. Map of site:

Refer to Annex III of the *Explanatory Note and Guidelines*, for detailed guidance on provision of suitable maps, including digital maps.

a) A map of the site, with clearly delineated boundaries, is included as:

- i) a **hard copy** (required for inclusion of site in the Ramsar List): ☐;
- ii) an **electronic format** (e.g. a JPEG or ArcView image) ☐;
- iii) a **GIS file providing geo-referenced site boundary vectors and attribute tables** ☐.

b) Describe briefly the type of boundary delineation applied:

e.g. the boundary is the same as an existing protected area (nature reserve, national park, etc.), or follows a catchment boundary, or follows a geopolitical boundary such as a local government jurisdiction, follows physical boundaries such as roads, follows the shoreline of a waterbody, etc.

8. Geographical coordinates (latitude/longitude, in degrees and minutes):

Provide the coordinates of the approximate centre of the site and/or the limits of the site. If the site is composed of more than one separate area, provide coordinates for each of these areas.

Latitude: 19° 27' S; Longitude: 147° 15' E

9. General location:

Include in which part of the country and which large administrative region(s) the site lies and the location of the nearest large town.

Centre 52km SE of the major provincial city of Townsville; 21km NE of the Township of Ayr.

10. Elevation: (in metres: average and/or maximum & minimum)

Mountainous areas: Mt Cleveland to 558m
 Feltham Cone to 279m
 The Cone to 94m

Remaining areas: All less than 20m; most less than 2m.

11. Area: (in hectares)

Total area 35 500ha divided as:

(a) Mountainous areas above 20m: Cape Cleveland, The Cone, Storth Hill, Feltham Cone: 4000ha or approximately 11% of the area.

(b) Elevated Coastal Sand Dunes above 10 meters: only small areas of this land-type fall within the proposed area;

(c) Coastal Plain below 20 meters, most much less: approximately 32 000ha or 89% of the total area.

12. General overview of the site:

Provide a short paragraph giving a summary description of the principal ecological characteristics and importance of the wetland.

The Ramsar site is a diverse complex of coastal wetland systems formed on four broad physiographic types viz., mountainous areas of Cape Cleveland and Feltham Cone; the elevated parallel dune systems; the lower lying parts of the coastal plain; and the actively prograding sand spit of Cape Bowling Green. Over most of the site, coastal mangrove communities give way inland to the highly saline communities of the salt pans which in turn lead to the brackish and freshwater communities of the lower lying coastal plain further inland. These lowland areas are typified by communities whose dominant ecological characteristic is a tolerance of saline conditions. Extensive areas of forest and woodland, and some closed forest, occur on the mountainous areas and the coastal dune system.

13. Ramsar Criteria:

Tick the box under each Criterion applied to the designation of the Ramsar site. See Annex II of the *Explanatory Notes and Guidelines* for the Criteria and guidelines for their application (adopted by Resolution VII.11). All Criteria which apply should be ticked.

1 •	2 •	3 •	4 •	5 •	6 •	7 •	8 •	9 •
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

14. Justification for the application of each Criterion listed in 13 above:

Provide justification for each Criterion in turn, clearly identifying to which Criterion the justification applies (see Annex II for guidance on acceptable forms of justification).

Criterion 1: A wetland should be considered internationally important if it contains a representative, rare, or unique example of a natural or near-natural wetland type found within the appropriate biogeographic region.

[Justification against former **Criterion 1(a)** under the Pre-1999 Criteria]:

The site is representative of many coastal and seasonal wetlands in the area, but is particularly significant for its diversity and extent of wetland types.

[Justification against former **Criterion 1(b)** under the Pre-1999 Criteria]:

The Bowling Green Bay site is representative of the richest coastal wetland habitats typical of the coastal wet-dry tropics of northeastern Australia.

[Justification against former **Criterion 1(c)** under the Pre-1999 Criteria]:

The site plays a substantial role in protection against erosion by cyclones. The site provides an extremely important bait fish, finfish and crustacean nursery, providing critical habitat at particular stages in their lifecycle.

Criterion 2: A wetland should be considered internationally important if it supports vulnerable, endangered, or critically endangered species or threatened ecological communities.

[Justification against former **Criterion 2(a)** under the Pre-1999 Criteria]:

The site provides feeding grounds for vulnerable species such as dugongs, estuarine crocodiles and green turtles, and several bird species.

Criterion 3: A wetland should be considered internationally important if it supports populations of plant and/or animal species important for maintaining the biological diversity of a particular biogeographic region.

[Justification against former **Criterion 2(b)** under the Pre-1999 Criteria]:

The site is particularly important for the abundance and diversity of avifauna and sustaining the genetic diversity of the region is dependent on maintenance of the site for continued use.

[Justification against former **Criterion 3(b)** under the Pre-1999 Criteria]:

The site regularly supports substantial numbers of all Australian waterbird groups, including post breeding populations of 4000 brolgas, magpie geese (10,000), and various species of Anatidae (4000-5000).

Criterion 4: A wetland should be considered internationally important if it supports plant and/or animal species at a critical stage in their life cycles, or provides refuge during adverse conditions.

[Justification against former **Criterion 2(c)** under the Pre-1999 Criteria]:

The site is of special significance as breeding and feeding habitat for brolgas and magpie geese.

Criterion 5: A wetland should be considered internationally important if it regularly supports 20,000 or more waterbirds.

[Justification against former **Criterion 3(a)** under the Pre-1999 Criteria]:

The site seasonally supports a wide range of waterbird species well in excess of 20,000 individuals.

Criterion 6: A wetland should be considered internationally important if it regularly supports 1% of the individuals in a population of one species or subspecies of waterbird.

[Justification against former **Criterion 3(c)** under the Pre-1999 Criteria]:

Although population sizes are poorly known in northern Australia, it is likely that, seasonally, populations of brolgas exceed 1% of the total population.

15. Biogeography (required when Criteria 1 and/or 3 and /or certain applications of Criterion 2 are applied to the designation):

Name the relevant biogeographic region that includes the Ramsar site, and identify the biogeographic regionalisation system that has been applied.

a) biogeographic region:

b) biogeographic regionalisation scheme (include reference citation):

16. Physical features of the site:

Describe, as appropriate, the geology, geomorphology; origins - natural or artificial; hydrology; soil type; water quality; water depth, water permanence; fluctuations in water level; tidal variations; downstream area; general climate, etc.

(a) Geology and geomorphology: The mountainous areas comprising Cape Cleveland, The Cone and Feltham Cone, occur in the northwestern part of the site. They are largely comprised of adamellite and granite of Permian to Mesozoic age, with small areas of older intermediate lavas and pyroclastics. Extensive areas of Quaternary sand dunes occur on the western side of Bowling Green Bay to the north of the Haughton River estuary. At the Eastern side of Bowling Green Bay, narrow sand dunes comprise much of the long fingerlike spit forming Cape Bowling Green. The remaining, most extensive area of the site comprises muds and silts of Quaternary age which, further inland, are replaced by the soils and sediments of the coastal plain.

(b) Origins: With the exception of the areas where water is impounded or impeded (indicated in 10 above), the wetlands of the site are the result of natural processes.

(c) Climate: The site is in a rain shadow area and lies between areas of substantially higher rainfall to both the north and the south. Annual average precipitation is 1101mm at Ayr, 1256mm at Giru and 1200mm at Cape Cleveland (Anon. 1970). The rainfall occurs predominantly in summer and 70% falls in the months January to March. There are few figures for pan evaporation available for the area; the average annual evaporation for Townsville, some 52km from the centre of the site, is 1664mm (Anon.1970). Average annual temperatures at Ayr, 21km from the centre of the site are: average mean 23.3°C, average minimum 17.9°C and average maximum 29.1°C. The site is subject to the effects of severe tropical cyclones and lows during the months November to April which bring destructive winds and intense rainfall. The return period is in the order of 30-50 years. The heavy storm rains of the summer wet season have a notable effect in seasonally reducing the salinities of the shallow inshore marine areas (Walker 1981) and the surface soils of the saltpans and mangrove areas (Pomeroy 1990).

(d) Hydrology: The site is drained seaward into Bowling Green Bay by the Haughton River, three larger (Barramundi, Barratta and Sheep Station Cks) and many small creeks. Two large creeks (Cocoa Ck and Alligator Ck) and a number of minor creeks discharge into Cleveland Bay at the north western end of the site; a few small creeks drain into Upstart Bay at its southern end. The mountainous area to the south of Cape Cleveland and Feltham Cone are drained by seasonally active creeks into Cleveland and Bowling Green Bays. McKenzie and Emmett Creeks drain seasonal runoff from the steep northeastern slopes of Mt Elliott into extensive sedge swamps situated in a shallow basin whose eastern edge is fringed by The Cone, Storth Hill and Feltham Cone. Since its catchment extends well into the hinterland of Bowling Green Bay, the Haughton River runs for 5-6 months of the year. During periods of extreme flooding it overflows northward directly into the Cromarty-Cleveland complex of swamps and floods out into Cleveland Bay. Similarly, seasonal floodwaters from the Burdekin river overflow northward to discharge into bowling Green Bay.

Groundwater is stored in two main underground aquifers which are recharged almost entirely by stream flows. The older aquifers are the deepest (15-45 meters) and extend from the slopes of Mt Elliott across the Haughton River to the Burdekin delta. The younger aquifers are shallower and limited in extent to the Haughton Riverbed and some recently abandoned channels. Groundwater is also extracted from aquifers adjacent to the site at Barratta Creek (Cepecha and Kaminskas 1973).

(e) Soils: A varied range of soils occur within the site and correspond largely to local elevation and the parent material present. The soils have been little studied in the southern and eastern parts of the site, although those to the northwest of the Haughton River have been described and mapped by Murtha (1982). The soils of the mountainous area associated with Cape Cleveland and Feltham Cone are derived largely from coarse-grained granitic parent materials. These are mostly thin sandy soils although occasional red soils with a marked texture contrast (duplex soils) also occur. On the piedmont slopes surrounding these areas, well drained earthy-textured soils (alfisols) occur.

Soils of the sand dune areas vary in degree and kind of profile development, depending on differences in age and possibly parent materials. Most are freely drained, coarse textured sandy soils with uniform texture profiles, although some weakly developed podzols with seasonal ponding of water occur in the swales.

The soils and sediments of the mangrove forests are poorly known and have been characterised by Murtha (1982) as largely "dark brown muds of unknown depth", although some sandy soils also support mangroves in the region. Limited analyses of these soils and sediments have been carried out within the site. Similarly, only limited information is available on the soils of the salt pans, mostly

from the above sources. Landwards from the mangrove forests of the coastal plain, sequences of soils reflect the influence of increasing elevation and changing salinity.

Surrounding the salt pans, and occasionally present as islands within them, are the duplex soils (alfisols) of the Coonambelah series. These are moderately to strongly gleyed and somewhat alkaline at depth and are occasionally flooded by the higher tides. Other soils of the coastal plain include the cracking clays (verisols) of the Brolga series found largely to the northwest of Feltham Cone. Situated slightly higher in the landscape are the alfisols of the Doughboy series which are duplex soils but which do not appear to be influenced by regular seawater inundation. All of the above soils evidence their often lengthy periods of summer inundation by the presence of gleying and many of the soils of the coastal plain reflect the influence of smectites in their propensity to crack on drying and swell on subsequent rehydration.

(f) Water Quality: There are few published water quality measure available for the area. The mountainous area of Cape Cleveland is mapped as having less than $1000 \mu\text{g g}^{-1}$ of total dissolved solids, while lower lying areas are mapped as having total dissolved solids in the range $1000\text{--}3000 \mu\text{g g}^{-1}$, or their being insufficient data. The latter areas include the salt pans, mangroves and other areas of known saline and brackish waters. It also includes the areas of coastal sand dunes at the western end of Bowling Green Bay. Aquifers in the hinterland of the site are of mixed quality, while waters of the younger alluvium are of good quality. Those of the older alluvium and those close to the coast may have total dissolved salts exceeding $3000 \mu\text{g g}^{-1}$. There is a risk of saltwater intrusion into the freshwater aquifers if excessive agricultural consumption occurs (Ceplecha and Kaminskas 1973).

Between the Haughton and Burdekin Rivers extensive areas of agricultural development are located upstream of all watercourses flowing into Bowling Green Bay. It is assumed that agricultural chemicals and nutrients are flushed into the area during periods of local flooding. The impact of this on the water quality of the site is unknown. However, it is likely to be significant since recharge of underground aquifers appears to be direct infiltration over the delta from rainfall, river flow and flood (Ceplecha and Kaminskas 1973). Water quality is least likely to be affected by agricultural contaminants for that portion of the site to the north of the Haughton River.

(g) Water depth, fluctuations and permanence: there is considerable variation throughout the site as indicated below.

- (i) Sub-tidal area - below mean low water neaps and permanently flooded with sea water.
- (ii) Riverine channels - varying depth and permanently flooded with sea water except during periods of wet season freshwater discharge when salinities fall to below 0.5 parts per thousand.
- (iii) Mangrove areas - shallow area (less than 2.0m) inundated diurnally to monthly, depending on tidal range.
- (iv) Saltpan areas - shallow areas (less than 0.8m) occasionally flooded by oceanic water and seasonally by freshwater.
- (v) Brackish areas - shallow areas (less than 2.0m) seasonally and semi-permanently flooded with sea water except during periods of wet season freshwater discharge when salinities fall to below 0.5 parts per thousand.
- (vi) Freshwater areas - shallow areas (less than 2.0m) seasonally inundated with water.

- (h) **Tidal variations:** The highest astronomical tides and mean high water springs are, respectively, 2.25 and 1.15m above mean sea level.
- (i) **Catchment area:** The major catchment is located beyond the extent of the site although there are some internal catchments associated with The Cone, Storth Hill, Feltham Cone and Cape Cleveland. Beyond the site, the Mt Elliott catchment immediately adjacent, and the catchments of Majors Creek, Haughton River and the Barratta Creek some 70km inland, drain the eastern slopes of the coastal ranges and some outliers. Land use within the catchment area is almost entirely agricultural and pastoral. The Burdekin river which drains some two thirds of Queensland overflows northward into Bowling Green Bay during periods of seasonal flooding.
- (j) **Downstream area:** No relevant comment.

17. Physical features of the catchment area:

Describe the surface area, general geology and geomorphological features, general soil types, and climate (including climate type).

18. Hydrological values:

Describe the functions and values of the wetland in groundwater recharge, flood control, sediment trapping, shoreline stabilization, etc.

- (a) **Recharge and discharge of ground water:** no information is available for the site or its role in the recharge and discharge of ground water. However, because of the extent of the fresh and brackish swamps it is likely to play a significant role in the recharge of underground fresh and saline aquifers from direct infiltration, rainfall, river flow and floods (Ceplecha and Kaminskas 1973).
- (b) **Sediment trapping:** The mangrove communities of Bowling Green Bay play an important role in the trapping of tide born sediments.
- (c) **Prevention of coastal erosion:** The mangrove communities exert a major influence in the control of coastal erosion in Bowling Green Bay. The area is subject to tropical cyclones and lows during the summer wet season. The strong winds, tidal surges and the intense and sustained rainfalls associated with these, all have considerable potential for eroding the foreshore and the coastal streams and rivers. Throughout the site mangrove and other wetland plant communities play a critical role in limiting the severity of erosive losses.
- (d) **Maintenance of water quality:** There is a delicate balance of salinity in the area, which has important implications for the seasonal distribution, dynamics and productivity of the vegetation of the fresh and brackish water swamps and of the tidal wetlands.
- (e) **Support of food chains:** Extensive bait fish breeding occurs in the site and supports important bill (black marlin, sail fish) and pelagic fisheries (spanish mackerel and other predatory species). In addition, the role of these wetlands in exporting carbon to offshore sites contributes to the overall productivity of the adjacent marine system. The intertidal and subtidal seagrass beds of both Bowling Green and Cleveland Bays directly provide feeding habitat for the threatened herbivores *Dugong dugon* and *Chelonia mydas* and nursery areas for a prawn industry. Commercial and recreational harvesting of prawns and the mud crab (*Scylla serrata*) with the site and adjacent marine areas is based on the productivity of the local wetland communities. The importance of the wetlands of the site and of Bowling Green Bay has been recognised in the protection of the entire bay as a Fish Habitat Area (gazetted 25 November 1989).

Further Inland the brackish and freshwater wetlands of the site provide an abundant source of food for a diverse and rich waterbird community: in particular the tubers of the dominant freshwater sedge

Eleocharis dulcis provide the major source of food for populations of the Brolga and Magpie goose which concentrate on these swamps in the post breeding season (Blackman 1978). In addition, while no published work is available, the fresh and brackish swamps also support an important local fishery based on the barramundi (*Lates calcarifer*). Tagging studies have shown that juveniles of this species breed prolifically in the fresh water swamps of the site and these are later found in the estuarine waters of Bowling Green Bay.

19. Wetland Types

a) presence:

Circle or underline the applicable codes for the wetland types of the Ramsar "Classification System for Wetland Type" present in the Ramsar site. Descriptions of each wetland type code are provided in Annex I of the *Explanatory Notes & Guidelines*.

Marine/coastal: A • B • C • D • E • F • G • H • I • J • K • Zk(a)

Inland: L • M • N • O • P • Q • R • Sp • Ss • Tp • Ts • U • Va •
Vt • W • Xf • Xp • Y • Zg • Zk(b)

Human-made: 1 • 2 • 3 • 4 • 5 • 6 • 7 • 8 • 9 • Zk(c)

b) dominance:

List the wetland types identified in a) above in order of their dominance (by area) in the Ramsar site, starting with the wetland type with the largest area.

20. General ecological features:

Provide further description, as appropriate, of the main habitats, vegetation types, plant and animal communities present in the Ramsar site, and the ecosystem services of the site and the benefits derived from them.

A range of different habitats integrating with one another occur throughout the site (Blackman 1978).

(a) Mountainous habitats: Open forest and woodland communities (*E. crebra*, *E. playtphylla*, *E. papuana*) predominate, with complex associations of vine thickets and *Arucaria cunninghamii* emergents in protected gullies.

(b) Coastal dune habitats: These comprise dune swale systems of predominantly *Casuarina equisetifolia* on the foredunes with Eucalypt and Melaleuca open forest and woodland (*E. tessellaris*, *E. tereticornis*, *E. alba*, *Lophostemon suaevoleus*, *Melaleuca leucodendron*, *M. quinquenervia*) on the older dunes behind these. Submerged and emergent aquatics, grasses, sedges, palms and a complex of littoral forest - often dominated by *Melaleuca* species - occur in the swales.

(c) Mangrove/tidal forest habitats: These comprise closed, open and woodland communities dominated by *Avicennia eucalyptifolia*, *Ceriops tagal* and *Rhizophora stylosa*. Within the site some nine species of mangrove occur, three of which (*Aegiceras corniculatum*, *Osbornia octodonta* and *Lumnitzera racemosa*) are poorly represented.

(d) Tidal mudflat habitats: On the seaward side these comprise open largely unvegetated mudflats. Inland, behind the fringing communities, the most hypersaline of the areas are devoid of macrophytic vegetation, while other less saline areas are dominated by open forblands comprising samphire communities. In some areas, causeways associated with roads have impeded water flow and impounded saline waters, resulting in an invasion of mangrove communities into these mudflat habitats.

(e) Salt Marsh habitats: These comprise slightly elevated plains dominated by *Sporobolus virginicus* and brackish sedge communities dominated by *Scirpus littoralis*.

(f) Stream and tidal channel habitats: These comprise open water and unconsolidated bottoms frequently with fringing mangrove, grass and samphire communities in the saline areas, while fringing forest and woodland communities dominated by *Melaleuca* species occur in brackish and freshwater areas.

(g) Freshwater marsh habitats: these comprise extensive sedge communities dominated by *Eleocharis dulcis* and a range of submerged and emergent aquatic communities, depending on the water regime. A number of areas within these habitats have been modified by the construction of earth walls which have increased the extent of freshwater habitat.

21. Noteworthy flora:

Provide additional information on particular species and why they are noteworthy (expanding as necessary on information provided in 14, Justification for the application of the Criteria) indicating, e.g., which species/communities are unique, rare, endangered or biogeographically important, etc. *Do not include here taxonomic lists of species present – these may be supplied as supplementary information to the RIS.*

The flora of the site is noteworthy because of the extensive wetland communities of intertidal algae, seagrasses, mangroves, saline grasses and herbs, sedges and other aquatic species, as well as the distinctive seasonal wetland communities of the forested swales. Similarly, the varied terrestrial communities present - open and closed Eucalypt and Melaleuca dominant communities of the beach ridges and the granitic hills, the monsoon and Araucarian closed forests of the sheltered slopes - provide a representation of the major coastal communities of the north Australian wet-dry tropics. Three species classified as rare or threatened by QPWS (*Cynanchum leptolepus*, *Bonamia dietrichiana*, *Livingstonia drudei*) occur within the site.

22. Noteworthy fauna:

Provide additional information on particular species and why they are noteworthy (expanding as necessary on information provided in 14, Justification for the application of the Criteria) indicating, e.g., which species/communities are unique, rare, endangered or biogeographically important, etc., including count data. *Do not include here taxonomic lists of species present – these may be supplied as supplementary information to the RIS.*

The avifaunal diversity of the site reflects the range of terrestrial and marine coastal habitats present. A provisional list of bird species recorded from within the site is provided as **Appendix 1**. Of the 224 birds known to occur in the site, 103 are known to breed within its confines; these are flagged in **Appendix 1** as are a total of 13 species that are rare or whose conservation status is endangered, vulnerable or rare (Draft list of the Conservation Status of Queensland Vertebrates, Queensland Parks and Wildlife Service, unpublished). Currently, most information is available for the bird fauna. Information on the other vertebrate species is too incomplete for inclusion at this time, although this is being redressed by surveys in progress. Little information is available on invertebrate fauna of the site.

The site is important habitat for some 50% of species listed in the Appendices of both the Japan Australia Migratory Bird Agreement (JAMBA - 38 of the 76 species with four (4) breeding) and the China Australia Migratory Bird Agreement (CAMBA - 40 of the 81 species with three (3) breeding). The site also provides habitat species listed in the Appendices of the Bonn Convention (two (2) endangered sea turtle species; two (2) species - estuarine crocodile and dugong - with unfavourable conservation status). **Appendix 2** provides a list of the above species.

One of the most noteworthy aspects of the avifauna of the site is the richness and local abundance of species present (Blackman 1978). This diversity is particularly evident within the waterbirds at both Family and Species levels - see the **Appendix 1** representatives of the Podicipitiformes, Pelecaniformes, Ardeiformes, Anseriformes, Gruiformes and Charadriiformes. This is also evident within other groups notably the Accipitriformes.

The brolga and magpie goose are the most noteworthy species of the shallow sedge swamp and marine plains of the site (Blackman 1988). Both species breed extensively in the sedge swamps during late summer. In addition, the site supports a post breeding (May - September) concentration of regional populations of both species. These concentrations are the largest in north east Queensland and the site forms part of the main stronghold of these two species in eastern Australia. Other notable species include the little tern which breeds on Bowling Green Bay spit and occurs in flocks of 1000 individuals; and dugongs and green turtles which depend on the extensive areas of intertidal sea grass beds.

23. Social and cultural values:

a) Describe if the site has any general social and/or cultural values e.g., fisheries production, forestry, religious importance, archaeological sites, social relations with the wetland, etc. Distinguish between historical/archaeological/religious significance and current socio-economic values:

The site is of a high aesthetic and cultural value. It has enormous potential for environmental education and nature based recreation, both of which have still to be realised.

b) Is the site considered of international importance for holding, in addition to relevant ecological values, examples of significant cultural values, whether material or non-material, linked to its origin, conservation and/or ecological functioning?

If Yes, tick the box ☐ and describe this importance under one or more of the following categories:

- i) sites which provide a model of wetland wise use, demonstrating the application of traditional knowledge and methods of management and use that maintain the ecological character of the wetland:
- ii) sites which have exceptional cultural traditions or records of former civilizations that have influenced the ecological character of the wetland:
- iii) sites where the ecological character of the wetland depends on the interaction with local communities or indigenous peoples:
- iv) sites where relevant non-material values such as sacred sites are present and their existence is strongly linked with the maintenance of the ecological character of the wetland:

24. Land tenure/ownership:

a) within the Ramsar site:

The greater portion of the site is gazetted as National Park, or proposed as such. The bulk of the site comprises a portion of Bowling Green Bay National Park which is under the management of the Queensland Parks and Wildlife Service. The site also includes a number of portions of Unallocated State Land, Leasehold land, and Special Purpose reserves all of which are designated for future inclusion into the National Park. The site surrounds, or is contiguous with portions of freehold and leasehold land. Below Mean High Water Springs, the site is contiguous with the Great Barrier Reef Marine Park and the Bowling Green Bay Fish Habitat Reserve which overlays the Marine Park in that area.

b) in the surrounding area:

Lands contiguous with the site comprise portions of privately owned freehold land; Commonwealth and Local Government land: and State land leased under a variety of tenures.

25. Current land (including water) use:

a) within the Ramsar site:

Livestock (cattle) grazing is carried out in portions of the site, mainly in the brackish and freshwater areas north west of the Haughton River. Grazing of these areas is crucial to the maintenance of the waterbird habitat.

b) in the surroundings/catchment:

The northern seaward portion of the site abuts the Great Barrier Reef Marine Park. Grazing is the predominant land use in almost all the adjacent portions continuous with the inland boundaries of the site. However, several small townships (Cungulla, Hucks Landing) are encompassed by the site but do not form part of it. At the northern extremities of Capes Bowling Green and Cleveland, small lighthouse reserves (approx. 80ha and 3ha respectively) occur and a further (208ha) has been excised at the western end of Bowling Green Bay for the Australian Institute of Marine Science. Extensive agricultural lands associated with the Burdekin river irrigation area occur further inland.

26. Factors (past, present or potential) adversely affecting the site's ecological character, including changes in land (including water) use and development projects:

a) within the Ramsar site:

The low lying topography of the site would make it extremely susceptible to marine flooding as the result of relatively small rises in sea level, one of the predicted outcomes of the greenhouse effect. Some simulation modeling has been undertaken to show the potential extent of this over the area of the site to the north of the Haughton River (Marshall and Blackman, in press).

b) in the surrounding area:

Agricultural, urban and industrial developments in the catchment area external to the site pose the most significant threats to the integrity of the site, particularly the lowland areas. The potential impact posed by these is through cumulative changes to water regimes and the chemistry of both surface and sub-surface waters. As already evident within the coastal region, such changes initiate rapid changes in biological communities and degrade the natural functioning of wetlands within the landscape and their value as habitat (Blackman and Brooke, in press).

27. Conservation measures taken:

a) List national and/or international category and legal status of protected areas, including boundary relationships with the Ramsar site:

In particular, if the site is partly or wholly a World Heritage Site and/or a UNESCO Biosphere Reserve, please give the names of the site under these designations.

The 32 100ha site is wholly within the Bowling Green Bay National Park (767) with the exceptions specified in 15 (a) above. The Park was established in 1977. Within the State of Queensland, National Parks are gazetted under the Nature Conservation Act 1992. National Park status of the site therefore represents one of the most stringent forms of protection available in Queensland.

b) If appropriate, list the IUCN (1994) protected areas category/ies which apply to the site (tick the box or boxes as appropriate):

Ia ☐; Ib ☐; II ☐; III ☐; IV ☐; V ☐; VI ☐

c) Does an officially approved management plan exist; and is it being implemented?:

No management plan is current for the site. However, Government policy requires that management plans be put into operation and a plan for Bowling Green Bay National Park is currently in preparation.

d) Describe any other current management practices:

Despite the lack of a general management plan, portions of the park are subject to control burning practices to reduce the damaging effects of wildfire. The shallow seasonal wetlands are also subject to controlled grazing to maintain the wetlands for waterfowl by preventing the invasion of upland grasses into the swamps during the dry spring. Limited public access is available to the whole park area for fishing and camping. Minor facilities are available at one location (Barramundi Ck) and proposed for another (Bald Rock, Cape Cleveland) The area is actively patrolled by two full time rangers of the QPWS who enforce regulations to the park.

28. Conservation measures proposed but not yet implemented:

e.g. management plan in preparation; official proposal as a legally protected area, etc.

Current surveys of the vegetation of the site are aimed at description and mapping of the vegetation communities and the production of species lists. Similarly, fauna surveys are planned to provide information on the distribution, abundance and habitat associations of mammals, birds, reptiles and frogs occurring in the Park.

29. Current scientific research and facilities:

e.g., details of current research projects, including biodiversity monitoring; existence of a field research station, etc.

Programs to map and characterise the natural resources (landscape, fauna, flora and the communities present) of the site are currently in progress.

30. Current communications, education, participation and awareness (CEPA) activities related to or benefiting the site:

e.g. visitors' centre, observation hides and nature trails, information booklets, facilities for school visits, etc.

No specific programs currently exist for the site.

31. Current recreation and tourism:

State if the wetland is used for recreation/tourism; indicate type(s) and their frequency/intensity.

The major recreational pursuits are fishing, camping, bushwalking, birdwatching and other nature related activities. At present there is no commercial exploitation of the site.

32. Jurisdiction:

Include territorial, e.g. state/region, and functional/sectoral, e.g. Dept of Agriculture/Dept. of Environment, etc.

QPWS has jurisdiction over the terrestrial areas of the site. It shares jurisdiction over the adjacent Bowling Green Bay Management Area of the Townsville - Whitsunday Marine Park. The Beach Protection Authority has some jurisdiction over the foreshores of the site and Queensland Department of Primary Industries has jurisdiction over the Fish Habitat Areas.

33. Management authority:

Provide the name and address of the local office(s) of the agency(ies) or organisation(s) directly responsible for managing the wetland. Wherever possible provide also the title and/or name of the person or persons in this office with responsibility for the wetland.

Queensland Parks and Wildlife Service
160 Ann Street
BRISBANE, QLD, 4000

34. Bibliographical references:

Scientific/technical references only. If biogeographic regionalisation scheme applied (see 15 above), list full reference citation for the scheme.

- Aliano, W.L (1978). *A study of the clay of an area on Cape Bowling Green*. Unpublished B.Sc. (Hons) thesis, Department of Geology, James Cook University of North Queensland.
- Anon. (1970). *Burdekin-Townsville Region - Climate*. Queensland Resources Series. Geographic Section, Department of National Development, Canberra. 19pp 1 map.
- Belperio, A.P (1978). *An inner shelf sedimentation model for the Townsville region, Great Barrier Reef Province*. Unpublished Ph.D thesis, Department of geology, James Cook University of North Queensland.
- Blackman, J.G (1978) The Swamps. In "*Exploration North*" H.J Lavery (ed.) Richmond Hill: Melbourne.
- Blackman, J.G and Brooke, G. (in press). Towards an ecologically based information system for coastal wetlands in Queensland. In "*Rural Queensland - A Sustainable Future*". Royal Society for Queensland. Proceedings of a Symposium, Brisbane, 23 November, 1989.
- Burdekin Project Committee. (1977). *Resources and Potential of the Burdekin River Basin, Queensland*. Australian Government Publishing Service, Canberra.
- Cepilecha, V.J and Kaminskas, A. (1973) *Burdekin-Townsville Region, Queensland Water Resources Series*, Division of National Mapping, Dept of minerals and Energy, Canberra.
- Christian, G.S, Paterson, S.J, Perry, R.A, Slatyer, R.O, Stewart, G.A and Traves, D.M (1953). *Survey of the Townsville - Bowen region, North Queensland, 1950*. CSIRO, Melbourne. 87pp, 8 plates, 1 map.
- Gregory, C.M (1969). *1:250,000 Geological Series - Explanatory Notes - Ayr, Queensland*. Bureau of Mineral Resources, Canberra. 14pp, 1 map.
- Hopley, D. (1970). *The geomorphology of the Burdekin Delta, North Queensland*. James Cook University of North Queensland, Department of Geography Monograph No:1.
- Hopley, D. and Murtha, G.G (1975). *The Quaternary deposits of the Townsville coastal plain*. James Cook University of north Queensland, Department of Geography Monograph No:8.
- Hubble, G.D and Thompson, C.H (1953). *The soils and landuse potential of the Lower Burdekin Valley, North Queensland*. CSIRO Australia, Division of Soils, Soils and Land Use Series. No.10.
- Marshall, V.L and Blackman, J.G (submitted). Application of GIS methods to aspects of wetland planning project in north Queensland. In "*Proceedings of 1990 GIS International Symposium*", Brisbane, 16-20 September 1990: 10pp.
- Marsh, H and Saalfeld, W.K (1990). The distribution and abundance of dugongs in the Great Barrier Reef Marine Park south of Cape Bedford. *Australian Wildlife Research*, 17, 511-524.
- Murtha, G.G (1982) *Soils and Land Use on the Southern Section of the Townsville Coastal Plain, North Queensland*. CSIRO, Australia, Division of Soils, Soils and Land Use Series, No: 59. 78pp, 1 map.

- Ngan, Y. and Price, I.R (1980). Distribution of intertidal benthic algae in the vicinity of Townsville, tropical Australia. *Australian Journal of Marine and Freshwater Research*, 31, 175 - 191.
- Pomeroy, A.B (1990) *A systems approach to the geochemistry, mineralogy and sedimentation of a coastal catchment and tidal flats in the dry tropics near Townsville, North Queensland, Australia*. Unpublished Ph.D thesis, Department of Civil and Systems Engineering, James Cook University of North Queensland.
- Reid, R.E and Baker, D.E (1984) *Soils of the Lower Burdekin River - Barratta Creek - Haughton River Area, North Queensland*. Queensland Department of Primary Industries, Agricultural Chemistry Branch, Technical Report No. 22.
- Robertson, A.I and Duke, N.C (1987) Mangroves as nursery sites: composition of fish and crustaceans in mangroves and other nearshore habitats in tropical Australia. *Marine Biology*, 96, 193-205
- Soil survey Staff (1975) *Soil Taxonomy: a basic system of soil classification for making and interpreting soil surveys*. U.S Department of Agriculture handbook No. 436. Government Printer, Washington, (DC).
- Spenceley, A.P (1976). Unvegetated saline tidal flats in North Queensland. *Journal of Tropical Geography*, 42, 78-85.
- Spenceley, A.P (1982). *The geomorphological and zonal development of mangrove swamps in the Townsville area, North Queensland*. Monograph Series No.11, Department of Geography, James Cook University of North Queensland.
- Thomas, M.B and McDonald, W.J.F (1989) *Rare and Threatened Plants of Queensland*. Second Edition. Department of Primary Industries, Queensland Government.
- Walker, T. (1991). Seasonal salinity variations in Cleveland Bay, Northern Queensland. *Australian Journal of Marine and Freshwater Research*, 32 143-149.
- Williams, D.M (1990) Significance of coastal resources to sailfish and juvenile black marlin in northeastern Australia: an on going research program. pp 21-28. In "*Planning the Future of Billfishes*" R. Stround (ed.) Proceedings of the 2nd international Billfish Symposium, Kailua-Kona, Hawaii. National Coalition for Marine Conservation, Savanna, Georgia.
- Williams, D.M and Cappa, M. (1990) Life histories of clupeids in north-eastern Australia: preliminary data. pp70 - 74. In "*Tuna Baitfish in the Indo-pacific Region*" S.J.M Blaber and Copland, J.W (eds) Proceedings of a workshop, Honiara, Solomon Islands, 11-13 Dec, 1989. ACIAR Proceedings No. 30.

Name and Address of Compiler:

J G Blackman and A V Spain
Department of environment and Heritage
Queensland National Parks and Wildlife Service
Northern Regional Centre
Marlow St
Pallarenda QLD 4810

Ph (077) 74 1411

Fx (077) 21 1737

Please return to: **Ramsar Convention Secretariat, Rue Mauverney 28, CH-1196 Gland, Switzerland**
Telephone: **+41 22 999 0170** • Fax: **+41 22 999 0169** • e-mail: **ramsar@ramsar.org**

Appendix 1. Provisional list of bird species recorded from within the Bowling Green Bay Ramsar site.

(Not present).

Appendix 2. Species at Bowling Green Bay Ramsar Site listed in the Japan Australia Migratory Bird Agreement (JAMBA), the China Australia Migratory Bird Agreement (CAMBA) and the Bonn Convention

(Not present).